



Dr. Techy says,
"Always follow up on abnormal test results."

FEBRUARY, 2014
BULLETIN 15
COMMON QUESTIONS
SERIES NO. 5

FOUR COMMON CLINICAL SITUATIONS AND ACTIONS TO TAKE

- BASED ON THIS 6 HORMONE PANEL:**
- CORTISOL
 - CORTICOTROPIN (ACTH)
 - PREGNENOLONE
 - DEHYDROEPIANDROSTERONE (DHEA)
 - PROGESTERONE
 - TESTOSTERONE

#1—NORMAL SERUM LEVELS



TREAT PAIN SYMPTOMATICALLY

#3—LOW SERUM LEVELS OF ONE OR MORE HORMONES



1-Increase medical analgesia if patient complains of inadequate pain control
2-Replace low hormones and retest in one month

#2—HIGH SERUM LEVELS OF ONE OR MORE HORMONES




1-increases medical analgesia.
2-Use opioids if necessary.

#4—OPIOID PATIENT WITH LOW SERUM LEVELS OF ONE OR MORE HORMONES



Replace hormones and retest every 2 to 3 months

Dr. Hormone says,
"High serum hormone levels represent uncontrolled pain. Low serum levels usually need temporary replacement."



Contact Information:
Tennant Foundation
334 S. Glendora Ave.
West Covina, CA 91740-3043
Ph: 626-919-7476
Fax: 626-919-7497
E-mail: veractinc@msn.com



Information Network